

An influence functional for ultrasoft QCD

F. Guerin⁺ and E. Iancu^{*}

⁺Institut Non Lineaire de Nice, 1361 route des Lucioles, 06560 Valbonne, France

^{*}Service de Physique Théorique, CE-Saclay, 91191 Gif sur Yvette, France

Abstract

A real-time path integral for ultrasoft QCD is formulated. It exhibits a Feynman's influence functional. The statistical properties of the theory and the gauge symmetry are explicit. The correspondence is established with the alternative version, where a noise term enters a transport equation.

Ultrasoft QCD is an effective theory for gluons at high temperature T ($g(T) \ll 1$) at distance and time scales larger than $(gT)^{-1}$, i.e. the scales T^{-1} and $(gT)^{-1}$ have been integrated out [1, 2]. The two dominant features are the coloured collective excitations of the gluons, and the important energy dissipation towards smaller distance scales. This last feature forbids an imaginary time formalism. In the real time formalism, an effective action has been found [3] which exhibits simple and interesting features. Here, the time has to run forwards and backwards along the time axis, A^+ is the field component that lives on the forward branch, and A^- lives on the backward branch. The appropriate basis for writing this action is

$$A^{(1)} = (A^+ + A^-)/2 \quad , \quad A^{(2)} = A^+ - A^- \quad , \quad (1)$$

known as the Keldysh basis [4]. The effective action for ultrasoft QCD reads

$$S(A^{(1)}, A^{(2)}) = S_0(A^{(1)}, A^{(2)}) + S_{\text{IF}}(A^{(1)}, A^{(2)}), \quad (2)$$

where S_0 is the Yang-Mills action in this basis:

$$\begin{aligned} S_0(A^{(1)}, A^{(2)}) &= - \int_X \frac{1}{4} [F_{\mu\nu}^+ F^{+\mu\nu} - F_{\mu\nu}^- F^{-\mu\nu}] \\ &= - \frac{1}{2} \int_X (D^{(1)\mu} A^{(2)\nu} - D^{(1)\nu} A^{(2)\mu})^a (F_{\mu\nu}^{(1)a} + g f^{abc} A_\mu^{(2)b} A_\nu^{(2)c} / 4) \end{aligned} \quad (3)$$

where $\int_X = \int_{-\infty}^{+\infty} dt \int d^3x$, a is a color index, $D_x^{(1)ab} = \delta^{ab} \partial_X + g f^{abc} A^{(1)c}$ is the covariant derivative built with the field $A^{(1)}$ in the adjoint representation. Furthermore:

$$\begin{aligned} S_{\text{IF}}(A^{(1)}, A^{(2)}) &= - \int_X A^{(2)\mu}(X) j_\mu(X; A^{(1)}) \\ &+ i \int_{X,Y} \frac{1}{2} A^{(2)\mu}(X) K_{\mu\nu}(X, Y; A^{(1)}) A^{(2)\nu}(Y) \quad , \end{aligned} \quad (4)$$

where the trace over the colour indices is implicitly understood. $j^\mu(X; A^{(1)})$ is the response function, i.e. the response of the plasma to a small external perturbation at the appropriate scale. In this nonlinear theory

$$j^\mu(X; A^{(1)}) = \Pi_{ret}^{\mu\nu} A_\nu^{(1)} + \frac{1}{2} \Gamma_{ret}^{\mu\nu\rho} A_\nu^{(1)} A_\rho^{(1)} + \dots \quad (5)$$

j_μ is expressed in terms of the (retarded) propagator along jagged paths in a background coloured field $A^{(1)}(X)$ [5, 2]

$$j_\mu^a(X; A^{(1)}) = m_D^2 \left[-g_{0\mu} A_0^{(1)a}(X) + i \int_{v,v'} \int d^4Y v_\mu G_{ret}^{ab}(X, Y; A^{(1)}; \mathbf{v}, \mathbf{v}') \partial_{Y_0} (v' \cdot A^{(1)b}(Y)) \right] \quad (6)$$

where $m_D^2 = g^2 N T^2 / 3$, $\int_v = \int d\Omega_v / 4\pi$, and G_{ret} is the retarded Green function defined as the solution to:

$$i (v \cdot D_x^{(1)} + \hat{C}) G_{ret}(X, Y; A^{(1)}; \mathbf{v}, \mathbf{v}') = \delta^{(4)}(X - Y) \delta_{S_2}(\mathbf{v} - \mathbf{v}') \quad (7)$$

$$\hat{C} G_{ret}(X, Y; A^{(1)}; \mathbf{v}, \mathbf{v}') = \int_{v''} C(\mathbf{v}, \mathbf{v}'') G_{ret}(X, Y; A^{(1)}; \mathbf{v}'', \mathbf{v}') \quad (8)$$

with the condition $G_{ret}(X, Y) = 0$ for $X_0 < Y_0$. $v \cdot D_x^{(1)}$ is the covariant drift operator in the space-time direction $v^\mu = (1, \mathbf{v})$ ($\mathbf{v}^2 = 1$). The collision operator $\hat{C}$ is local in X , and changes the $\mathbf{v}$ direction of the drift. It is symmetric in $\mathbf{v}$ and $\mathbf{v}'$, and has positive eigenvalues of order $g^2 T \ln 1/g$, except for a zero mode $\int_v C(\mathbf{v}, \mathbf{v}') = 0$. Its explicit form can be found in Refs. [1, 5, 6, 2].

The dissipation comes from $\hat{C}$ and it gives rise to the second term in the action. $K_{\mu\nu}$ was found to be the complement to j_μ that is needed to fulfill the KMS constraints, which exist, for each n , on n -point amplitudes[3].

$$K_{\mu\nu}^{ab}(X, Y; A^{(1)}) = i m_D^2 T \int_{v,v'} \left[v_\mu G_{ret}^{ab}(X, Y; A^{(1)}; \mathbf{v}, \mathbf{v}') v'_\nu + v_\nu G_{ret}^{ba}(Y, X; A^{(1)}; \mathbf{v}, \mathbf{v}') v'_\mu \right]. \quad (9)$$

Note that $i G_{ret}(X, Y; A; \mathbf{v}, \mathbf{v}')$ is real, therefore $j_\mu(X; A^{(1)})$ is real, while $K^{\mu\nu}(X, Y; A^{(1)})$ is real and positive. For example

$$K^{\mu\nu}(p_0, \mathbf{p}; A^{(1)} = 0) = -\frac{T}{p_0} 2 \text{Im} \Pi_{ret}^{\mu\nu}(p_0, \mathbf{p}). \quad (10)$$

The occurrence of $K^{\mu\nu}(X, Y; A^{(1)})$ in the action may be interpreted as the extension to a nonlinear case of the fluctuation-dissipation relation.

As early as 1963, Feynman[7] introduced the influence functional $\exp iS_{\text{IF}}$. In a path integral approach, this factor accounts for the influence of the environment upon the system. Feynman examined the case of gaussian influence functionals, i.e. S_{IF} quadratic both in A^+ and A^- , and he showed [7] that the term linear in $A^{(2)}$ should be real, while the term

quadratic in $A^{(2)}$ should be imaginary. The action (4) exhibits these properties for the $A^{(2)}$ dependence, while it is fully non-linear in the non-abelian gauge field $A^{(1)}$. From Feynman's point of view, the modes of the QCD plasma with momenta $p \ll gT$ made up the “system”, while the role of the “environment” is played by the modes that have been integrated out, i.e., by the modes with momenta $p \geq gT$. This interpretation will be fully developed somewhere else [8]. Recent applications of the influence functional [9, 10] did not consider gauge theories.

Ultrasoft QCD is in fact a semi-classical theory [1, 2]. Indeed the number of gluons is $N_g(p_0 \ll T) \sim T/p_0 \gg 1$, i.e. $N_g \approx N_g + 1$, and the commutation relations' effect should be negligible. $A^{(1)}$ may be identified with a classical field, while $A^{(2)}$ describes the thermal fluctuations around it. The physical role of the $A^{(2)}$ field is to balance, via its fluctuation, the dissipation associated with the collision term, in such a way that the field $A^{(1)}$ remains in thermal equilibrium.

The action (3)–(4) is invariant under the gauge transformations of the $A^{(1)}$ field, with $A^{(2)}$ transforming covariantly, as well as j^μ and $K^{\mu\nu}$,

$$A^{(1)\mu} \rightarrow h A^{(1)\mu} h^\dagger - (i/g) h \partial^\mu h^\dagger, \quad A^{(2)\mu} \rightarrow h A^{(2)\mu} h^\dagger. \quad (11)$$

Moreover j_μ and $K^{\mu\nu}$ are covariantly conserved:

$$\begin{aligned} D_\mu^{(1)} j^\mu &= 0, \\ D_{x\mu}^{(1)} K^{\mu\nu} &= D_{y\nu}^{(1)} K^{\mu\nu} = 0. \end{aligned} \quad (12)$$

The generating functional of the n -point correlation functions of the fields $A^{(1)}, A^{(2)}$ at temperature T is the following path-integral [8]

$$Z(J_1, J_2) = \int \mathcal{D}A^{(1)} \mathcal{D}A^{(2)} \exp i[S(A^{(1)}, A^{(2)}) + J_2 A^{(1)} + J_1 A^{(2)}] \quad (g.f.) \quad (13)$$

where $(g.f.)$ is the gauge-fixing factor. Note that this is written fully in real time: there is no need for an imaginary-time piece in the contour since the thermalisation of the fields $A^{(1)}$ proceeds exclusively via their scattering off the thermal fluctuations of the “medium” (the hard modes integrated over, and simulated here by the fields $A^{(2)}$) [8].

The measure in the path integral (13) is invariant under the transformation (11). The normalization of $Z(J_1, J_2)$ is $Z(J_1, J_2 = 0) = \text{a constant}$. Indeed, this condition means that, in addition to $S(A^{(1)}, A^{(2)} = 0) = 0$, all the correlation functions, involving the field $A^{(2)}$ only, vanish, and this property is a characteristic feature of the Keldysh basis [4].

Moreover, the fields $A^{(2)}$ are typically weak, so their mutual interactions can be safely neglected [8]. Indeed, their fluctuations are suppressed by the large imaginary term in the action, which implies $\overline{A^{(2)}} \equiv \sqrt{\langle A^{(2)} A^{(2)} \rangle} = \sqrt{K^{-1}}$, with $K^{-1} \propto p_0/T \ll 1$. It is therefore appropriate to also neglect the term cubic in $A^{(2)}$ in the Yang-Mills action (3). Once this is done, the resulting full action has the simplifying feature to be quadratic in the “fluctuation” field $A^{(2)}$ (but non-linear to all orders in the “background field” $A^{(1)}$). Specifically :

$$\begin{aligned} S(A^{(1)}, A^{(2)}) &= \int_X A^{(2)\mu}(X) \left(D^{(1)\nu} F_{\nu\mu}^{(1)} - j_\mu(X; A^{(1)}) \right) \\ &\quad + i \int_{X,Y} \frac{1}{2} A^{(2)\mu}(X) K_{\mu\nu}(X, Y; A^{(1)}) A^{(2)\nu}(Y). \end{aligned} \quad (14)$$

This action is manifestly invariant under the gauge transformations (11), and it has the additional gauge symmetry $A_\mu^{(1)} \rightarrow A_\mu^{(1)}$, $A_\mu^{(2)} \rightarrow A_\mu^{(2)} + D_\mu^{(1)} \Lambda(X)$. One may then fix the gauge and perform the path integral on the $A^{(2)}$ field. The choice of a background-gauge fixing term for $A^{(2)}$, i.e. $i(D_\mu^{(1)} A^{(2)\mu})^2/2\lambda$, has the advantage that it preserves the gauge symmetry of $A^{(1)}$, while it introduces $\tilde{K}_{\mu\nu} = K_{\mu\nu} + D_\mu^{(1)} D_\nu^{(1)}/\lambda$. (An alternative gauge choice is the time-axial gauge $A_0^{(1)} = 0 = A_0^{(2)}$). As a result of the path integration on $A^{(2)}$, a *real* gaussian weight appears for the quantity $D^{(1)\nu} F_{\nu\mu}^{(1)} - j_\mu(X; A^{(1)})$, which represents the fluctuations around the classical equation of motion for the field $A^{(1)}$. The generating functional of the correlation functions of the field $A^{(1)}$ is now written as a path integral on the field configurations, with a real weight, as in a statistical mechanics approach.

The comparison with the alternative approach to ultrasoft QCD (which is in term of two classical equations) makes use of the identity

$$\exp(-A^{(2)} K A^{(2)}/2) = [\det K]^{-1/2} \int \mathcal{D}\eta \exp(-\eta K^{-1} \eta/2 - i\eta A^{(2)}) \quad (15)$$

where $A^{(2)} K A^{(2)} = A_i^{(2)} K_{ij} A_j^{(2)}$, $\mathcal{D}\eta = \mathcal{D}\eta_i(X)$, with the gauge choice $A_0^{(2)} = A_0^{(1)} = 0$. The action $S(A^{(1)}, A^{(2)}, \eta)$ is now linear in $A^{(2)}$, and the result of the path integration on $A^{(2)}$ is a functional δ -function which enforces an effective equation of motion for the field $A^{(1)}$:

$$D^{(1)\nu} F_{\nu i}^{(1)} - j_i(X; A^{(1)}) - \eta_i(X) = 0. \quad (16)$$

The current $\eta_i(X)$ is a random variable with, for *fixed* $A^{(1)}$, a real Gaussian distribution with zero average and 2-point correlation function:

$$\langle \eta_i^a(X) \eta_j^b(Y) \rangle|_{A^{(1)} \text{ fixed}} = K_{ij}^{ab}(X, Y; A^{(1)}). \quad (17)$$

In reality, however, $A^{(1)}$ is related to η via the non-linear equation of motion (16), so that the true statistics of the coupled random fields $A^{(1)}$ and η is considerably more involved. Moreover, in the gauge $A_0^{(2)} = A_0^{(1)} = 0$, the Coulomb law $D^{(1)\nu} F_{\nu 0}^{(1)} - j_0(X; A^{(1)}) - \eta_0(X) = 0$ is a consequence of Eqs. (16)–(12), with η_0 defined by $\partial_t \eta_0 + D_i \eta_i = 0$, and with vanishing initial conditions at $t = -\infty$. Eq. (17) is then generalized to

$$\langle \eta_\mu^a(X) \eta_\nu^b(Y) \rangle|_{A^{(1)} \text{ fixed}} = K_{\mu\nu}^{ab}(X, Y; A^{(1)}), \quad (18)$$

i.e., for *fixed* $A^{(1)}$, $K_{\mu\nu}$ is the 2-point correlation function of the random source η_μ .

The original formulation of the effective theory by Bödeker [1] involves a set of coupled equations

$$(D^\nu F_{\nu\mu})_a(X) = m_D^2 \int_v v_\mu W_a(X, \mathbf{v}) \quad (19)$$

$$(v \cdot D + \hat{C})^{ab} W_b(X, \mathbf{v}) = \mathbf{v} \cdot \mathbf{E}^a + \xi^a(X, \mathbf{v}) \quad (20)$$

where $\xi(X, \mathbf{v})$ is a gaussian white noise term, with zero average and the 2-point correlation function

$$\langle \langle \xi^a(X, \mathbf{v}) \xi^b(Y, \mathbf{v}') \rangle \rangle = \frac{2T}{m_D^2} C(\mathbf{v}, \mathbf{v}') \delta^{(4)}(X - X') \delta^{ab}. \quad (21)$$

where $\langle\langle \cdots \rangle\rangle$ means average over the random white noise. Eq. (20) may be solved with the help of the retarded Green function (7). With the initial condition $W = 0$ at $t = -\infty$, the result is

$$m_D^2 \int_v v_\mu W(X, \mathbf{v}) = j_\mu(X; A) + j_\mu^{(\xi)}(X; A) , \quad (22)$$

where $j_\mu(X; A) \equiv j_\mu(X; A^{(1)} = A)$ defined in Eq. (6) (with $\mathbf{v} \cdot \mathbf{E} = \partial^0(v \cdot A) - v \cdot D A^0$), and

$$j_\mu^{(\xi)}(X; A) = m_D^2 \int d^4 Z \int_{v, v'} i v_\mu G_{ret}(X, Z; A; \mathbf{v}, \mathbf{v}') \xi(Z, \mathbf{v}'). \quad (23)$$

Thus, Eq. (19) can be rewritten as:

$$D^\nu F_{\nu\mu}(X) = j_\mu(X; A) + j_\mu^{(\xi)}(X; A) , \quad (24)$$

where the second piece $j_\mu^{(\xi)}$ of the current is proportional to the random source ξ , and so is itself a random quantity. One notes that, for *fixed* A , $j_\mu^{(\xi)}$ has a Gaussian distribution with zero average and 2-point correlation function which follows from Eq. (21):

$$\langle\langle j_\mu^{(\xi)a}(X; A) j_\nu^{(\xi)b}(Y; A) \rangle\rangle|_{A \text{ fixed}} = K_{\mu\nu}(X, Y; A) , \quad (25)$$

with the use of the identity

$$\hat{G}^T 2\hat{C} \hat{G} = -i \hat{G}^T \left((\hat{G}^T)^{-1} + (\hat{G})^{-1} \right) \hat{G} = -i(\hat{G} + \hat{G}^T) , \quad (26)$$

where $\hat{G}^T$ is the transpose of $\hat{G}$ in X space, colour space, and $\mathbf{v}$ space, i.e. $i \hat{G}^T = (-v \cdot D + \hat{C})^{-1}$ in the adjoint representation.

Comparing Eqs. (24)–(25) to Eqs. (16)–(18), one concludes that the random current $j_\mu^{(\xi)}(X; A)$ can be identified with the variable $\eta_\mu(X)$, and the field $A(X)$ with $A^{(1)}(X)$. An attempt to write down a path integral, involving the field $A(X)$ only, was made by Arnold [11] in the case of the static approximation to the W field equation (20), and in the $A_0 = 0$ gauge.

An explicit comparison between the two approaches is possible when the Green function $G_{ret}(X, Y; A)$ is expanded in powers of A , an expansion valid for $P \sim g^2 T \ln 1/g$. The solution to the classical equations (19)(20) is then obtained as an expansion in powers of the noise ξ , while the expansion of S_{IF} (see Eq. (4)) in powers of $A^{(1)}$ produces effective propagators and 1PI vertices involving $A^{(1)}$ and $A^{(2)}$. It has been checked that both approaches give the same answer for the n -point correlation functions of $A^{(1)} \equiv A$: (i) at the tree-level for $n = 2, 3$ [3] and $n = 4$, (ii) at the one-loop level for $n = 2$, as it is now detailed. In the field theory approach, the result for $\langle A^{(1)}(X) A^{(1)}(Y) \rangle$ is, in momentum space (disregarding the indices μ, ν)

$$G_c(P) = \frac{T}{p_0} \left(\frac{1}{P^2 - \Pi_r(P) - \Pi_{r \text{ new}}(P)} - [P \rightarrow -P] \right) \quad (27)$$

where Π_r is the ultrasoft self-energy (see Eq.(5)) and $\Pi_{r \text{ new}}$ is the contribution to the self-energy arising from one-loop diagrams with loop-momenta $K \sim g^2 T \ln 1/g$ and with effective

vertices and propagators [12]. In the classical form $\langle\langle A(X)A(Y)\rangle\rangle$, the terms quadratic and quartic in the noise ξ lead to the two lowest-order terms in an expansion of $\Pi_{r\text{ new}}$ of the right hand side of Eq. (27).

To conclude, this effective theory provides a full explicit example of a Feynman influence functional for a nonlinear case [8], in contrast to other examples where the emphasis is on the “backreaction”. Note finally that a similar influence action can be constructed [8] for the HTL effective theory[2], i.e., the theory for the collective dynamics over the scale $(gT)^{-1}$.

References

- [1] D. Bödeker, *Phys. Lett.* **B426** (1998) 351; *Nucl. Phys.* **B559** (1999) 502.
- [2] J. P. Blaizot and E. Iancu, *Phys. Rep.* **359** (2002) 355, hep-ph/0101103.
- [3] F. Guerin, *Phys. Lett.* **B530** (2002) 142 ; *Nucl. Phys.* **B629** (2002) 233.
- [4] K. Chou, Z. Su, B. Hao, and L. Yu, *Phys. Rep.* **118** (1985) 1.
- [5] J. P. Blaizot and E. Iancu, *Nucl. Phys.* **B570** (2000) 326; **B557** (1999) 183.
- [6] P. Arnold, D. Son and L.G. Yaffe, *Phys. Rev.* **D59** (1999) 105020; **D60** (1999) 025007.
- [7] R.P. Feynman and A.R. Hibbs, *Quantum Mechanics and Path Integrals* (McGraw-Hill, NewYork, 1965).
- [8] F. Guerin and E. Iancu, in preparation.
- [9] C. Greiner and B. Muller, *Phys. Rev.* **D53** (1997) 1026.
- [10] B.L. Hu, p.309, in *Banff/CAP workshop on Thermal Field Theory* (World Scientific, Singapore, 1994).
- [11] P. Arnold, *Phys. Rev.* **D62** (2000) 036003.
- [12] F. Guerin, *Phys.Rev.* **D63** (2001) 045017.